

Log-aesthetic curves and generalized Archimedean spirals

Péter Salvi

Budapest University of Technology and Economics

CGTA'25

Sopron, June 16–20, 2025

Outline

Introduction

- Classical Aesthetic Curves
- Logarithmic Curvature Histogram
- Log-Aesthetic Curves
- Generalized Archimedean Spirals

The Connection

- Radial Curves

Approximation

- LCH Slope of GA spirals
- Two Methods



Reconstruction

- Generalized Log-Aesthetic Curves

Conclusion





Classical Aesthetic Curves



Spline energies [Hoschek–Lasser '96]

$$\kappa''(s) = 0 \quad (\text{wooden}) \quad \Rightarrow \text{clothoid}$$

$$\int \kappa(s)^2 \, ds \rightarrow \min \quad (\text{mechanical}) \quad \Rightarrow \text{elastica}$$



Logarithmic Curvature Histogram (LCH)

Curve shape evaluation

[Harada et al. '99]:



1. Take samples of the curvature radius (ρ_i) at equal arc lengths
2. Divide $\ln(\rho_i)$ into a fixed number of bins
3. Plot the logarithm of the percentage of samples in the bins

→ : $\ln \rho$

$$\uparrow : \ln \frac{\partial s}{\partial \ln \rho} = \ln \frac{\partial s}{\partial \rho / \rho}$$

Straight lines are favorable

Logarithmic spiral



LCH—Alternative Interpretation


[Yoshida–Saito '06]:

1. Divide the curve into segments with the same $\Delta\rho/\rho$ ratio
2. Draw the log–log plot of segment lengths, i.e., $\ln(\Delta s)$ over $\ln(\rho)$

Linearity means

$$\kappa(s) = (c_0 s + c_1)^{-1/\alpha}$$

where α is the slope





Log-Aesthetic Curves [Miura '06]

$$\kappa(s) = (c_0 s + c_1)^{-1/\alpha}$$

$$\theta(s) = \frac{\alpha(c_0 s + c_1)^{(\alpha-1)/\alpha}}{(\alpha-1)c_0} + c_2$$

$$\mathbf{C}(s) = \mathbf{P}_0 + \left(\int_0^s \cos \theta(s) ds, \int_0^s \sin \theta(s) ds \right)$$



Types of Log-Aesthetic Curves


- ▶ Circle ($c_0 = 0$)
- ▶ Circle involute ($\alpha = 2$)
- ▶ Logarithmic spiral ($\alpha = 1$)
 - ▶ $\theta(s) = \ln(c_0 s + c_1)/c_0 + c_2$
- ▶ Nielsen's spiral ($\alpha = 0$)
 - ▶ $\kappa(s) = \exp(c_0 s + c_1)$
 - ▶ $\theta(s) = \exp(c_0 s + c_1)/c_0 + c_2$
- ▶ Clothoid ($\alpha = -1$)




Generalized Archimedean Spirals

Polar equation: $r = a + b\phi^{\frac{1}{c}}$

- ▶ $c = -2$: lituus
- ▶ $c = -1$: hyperbolic spiral
- ▶ $c = 1$: Archimedean (arithmetic) spiral
- ▶ $c = 2$: Fermat's spiral



Hyperbolic spiral



Fermat's spiral

Radial Curves


- ▶ Vector to the center of curvature, placed at the origin
- ▶ $\theta(t)$: tangent angle to the x axis
- ▶ $\theta^\perp(t) = \theta(t) + \frac{\pi}{2}$
- ▶ $\mathbf{R}(t) = [\cos \theta^\perp(t), \sin \theta^\perp(t)] \cdot \rho(t)$
- ▶ For log-aesthetic curves:

$$\rho(\theta^\perp) = \left(\theta^\perp c_0 \frac{\alpha - 1}{\alpha} \right)^{\frac{1}{\alpha - 1}}$$

- ▶ Polar equation:

$$r = b\phi^{\frac{1}{\alpha - 1}}$$

- ▶ GA spiral with $a = 0$ and $c = \alpha - 1$




Logarithmic spiral
(special case: $r = e^{b\phi}$)

LCH Slope of GA spirals


$$\alpha(t) = 1 + \frac{\rho(t)}{\rho'(t)^2} \left(\frac{\rho'(t)s''(t)}{s'(t)} - \rho''(t) \right)$$

Approaches $c + 1$ (slope of the related LA curve)



$c = -1$ (Hyperbolic spiral)

$\rightarrow \alpha = 0$ (Nielsen's spiral)



$c = 2$ (Fermat's spiral)

$\rightarrow \alpha = 3$

Approximating LA curves by GA spirals

- ▶ LA curve segment:


$$\mathbf{C}(s) = \mathbf{P}_0 + \left(\int_0^s \cos \theta(s) ds, \int_0^s \sin \theta(s) ds \right), \quad s \in [s_{\min}, s_{\max}]$$

- ▶ GA spiral segment:


$$\mathbf{C}_{\text{GA}}(t) = [\cos t, \sin t] \cdot (a + bt^{\frac{1}{c}}), \quad t \in [t_{\min}, t_{\max}]$$

- ▶ $a = 0$ and $c = \alpha - 1$
- ▶ Assume matching starting point and direction
 - ▶ Simple translation/rotation
- ▶ Interpolate curvature at $t_{\min}$
 - ▶ If $t_{\min}$ is known $\rightarrow b$ can be computed
- ▶ Interpolate curvature derivative at $t_{\min}$
 - ▶ $t_{\min}$ found by binary search
 - ▶ Initial frame by iterative doubling


Example 1: clothoid vs. lituus ($\alpha = -1$)



Example 2: Nielsen's spiral vs. hyperbolic spiral ($\alpha = 0$)



Example 3: Circle involute vs. arithmetic spiral ($\alpha = 2$)




Alternative Constraint

- ▶ Idea: Fix the endpoint instead of the curvature derivative
- ▶ Different error function for the bisection search
 - ▶ Radial distance of the endpoint to the GA spiral


Algorithm

1. Rotate the spiral s.t. $\mathbf{C}'_{\text{GA}}(t_{\min})$ points to $\theta(s_{\min})$.
2. Set $\mathbf{Q}$ (the spiral center) s.t. $\mathbf{Q} + \mathbf{C}_{\text{GA}}(t_{\min}) = \mathbf{P}_0$.
3. Let $\mathbf{u}$ and $\mathbf{v}$ be unit vectors from $\mathbf{Q}$ to $\mathbf{P}_0$ and $\mathbf{C}(s_{\max})$.
4. Set $t_{\max} = t_{\min} + \arccos\langle \mathbf{u}, \mathbf{v} \rangle$, or, if $\det(\mathbf{u}, \mathbf{v}) < 0$, choose the larger angle: $t_{\max} = t_{\min} + 2\pi - \arccos\langle \mathbf{u}, \mathbf{v} \rangle$.
5. The error is $\|\mathbf{C}(s_{\max}) - \mathbf{Q}\| - \|\mathbf{C}_{\text{GA}}(t_{\max})\|$.


Example 1b: clothoid vs. lituus ($\alpha = -1$)




Example 2b: **Nielsen's spiral** vs. hyperbolic spiral ($\alpha = 0$)




Example 3b: Circle involute vs. arithmetic spiral ($\alpha = 2$)



Example 4: Good Approximation ($\alpha = -\frac{3}{2}$, $t_{\min} \approx 9.88$)



Example 5: Bad Approximation ($\alpha = -1$, $t_{\min} \approx 1.42$)



Reconstructing Log-Aesthetic Curves from Radials

- ▶ From the construction: $\|\mathbf{C}'(t)\| = \|\mathbf{R}(t)\|$
- ▶ Inverse radial:

$$\mathbf{C}(t) = \int_0^t \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \cdot \mathbf{R}(t) dt$$

- ▶ Explicit equations for some cases, e.g.:
 - ▶ $b = 1, c = 1$ (circle involute):

$$[t \cos t - \sin t, t \sin t + \cos t]$$

- ▶ $b = 1, c = \frac{1}{2}$:

$$[(t^2 - 2) \cos t - 2t \sin t, (t^2 - 2) \sin t + 2t \cos t]$$


- ▶ etc.

- ▶ May involve incomplete gamma functions


Generalized Log-Aesthetic Curves

What if $a \neq 0$?

- ▶ Arithmetic spirals ($c = 1$): just a shift
- ▶ $c < 0$: LCH slope diverges into $\pm\infty$
- ▶ $c > 0$: Still converges to $c + 1$




$$c = \frac{1}{2}, a \in \{0, 20, 40, 60, 80\}$$



$$c = 2, a \in \{0, 1, 2, 3, 4\}$$

Conclusion

- ▶ Radial of LA curve $\rightarrow$ GA spiral
- ▶ LCH slope of radials approach α
- ▶ GA spirals approximate LA curves
- ▶ Reconstruction from radials
- ▶ Generalized LA curves



← **Elastica** vs. trig-aesthetic curves

References

- 1996 J. Hoschek, D. Lasser:
Fundamentals of Computer Aided Geometric Design. A. K. Peters.
- 1999 T. Harada, F. Yoshimoto, M. Moriyama:
An aesthetic curve in the field of industrial design.
Proceedings of IEEE Symposium on Visual Language, pp. 38–47.
- 2006 K. T. Miura: *A general equation of aesthetic curves and its self-affinity.*
Computer-Aided Design and Applications, Vol. 3, No. 1–4, pp. 457–464.
- 2006 N. Yoshida, T. Saito: *Interactive aesthetic curve segments.*
The Visual Computer, Vol. 22, pp. 896–905.
- 2024 N. Yoshida, T. Saito: *Shape information of curves and its visualization using two-tone pseudo coloring.*
Computer-Aided Design and Applications, Vol. 21, pp. 11–28.



<https://3dgeo.iit.bme.hu/>